



**PCRBIO SYSTEMS**  
simplifying research

## PCRBIO HS Taq Mix

[www.pcrbio.com](http://www.pcrbio.com)

### Product description

PCRBIO HS Taq Mix uses the latest developments in polymerase technology and buffer chemistry to enhance PCR speed, yield and specificity. The enzyme uses advanced hot start technology for superior sensitivity.

The enzyme and buffer system allow for superior PCR performance on complex templates such as mammalian genomic DNA. Due to enhanced efficiency and specificity the enzyme is perfectly suited to difficult PCR.

The mix contains PCRBIO HS Taq DNA Polymerase, a robust enzyme ideal for all your everyday PCR applications including genotyping, screening, library construction and multiplex PCR. The enzyme is particularly resistant to PCR inhibitors and is suitable for direct PCR from unprocessed samples including bacterial culture, bacterial colonies, blood and urine.

PCRBIO HS Taq Mix has an error rate of approximately 1 error per  $2.0 \times 10^5$  nucleotides incorporated. PCR products generated with PCRBIO HS Taq DNA Polymerase are A-tailed and may be cloned into TA cloning vectors.

High throughput screening has resulted in a buffer system that allows efficient amplification from GC-rich and AT-rich templates, under both fast and standard cycling conditions.

Component	200 reactions	1000 reactions
2x PCRBIO HS Taq Mix	5 x 1mL	25 x 1mL

### Shipping and storage

On arrival the kit should be stored between -30°C and -15°C. If stored correctly the kit will retain full activity for 12 months. The kit can be stored at 4°C for 1 month.

### Limitations of product use

The product may be used for in vitro research purposes only.

### Technical support

Help and support is available on our website at <https://pcrbio.com/resources/> including answers to frequently asked technical questions. For technical support and troubleshooting you can submit a technical enquiry online, or alternatively email [technical@pcrbio.com](mailto:technical@pcrbio.com) with the following information:

- Amplicon size
- Reaction setup
- Cycling conditions
- Screen grabs of gel images

## Important considerations

**2x PCRBIO HS Taq Mix:** The 2x mix contains PCRBIO HS Taq DNA Polymerase, 6mM MgCl<sub>2</sub>, 2mM dNTPs, enhancers and stabilizers. It is not recommended to add further PCR enhancers or MgCl<sub>2</sub> to the reaction. The buffer composition has been optimised to maximise PCR success rates.

**Template:** For eukaryotic DNA use between 5ng and 500ng per reaction, for cDNA use below 100ng per reaction.

**Primers:** Primers should have a predicted melting temperature of around 60°C, using default Primer 3 settings (<http://bioinfo.ut.ee/primer3/>). The final primer concentration in the reaction should be between 0.2µM and 0.6µM.

**Annealing:** We recommend performing a temperature gradient to experimentally determine the optimal annealing temperature. Alternatively, we recommend a 55°C annealing temperature then increase in 2°C increments if non-specific products are present.

**Extension:** Optimal extension is achieved at 72°C. The optimal extension time is dependent on amplicon length and complexity of template. 20 seconds per kilobase (kb) is recommended for amplification from eukaryotic DNA for amplicons between 1kb and 6kb. For shorter amplicons, faster cycling is possible.

**Multiplex PCR:** When first performing multiplex PCR it is recommended to run an annealing temperature gradient from 55°C to 65°C. The annealing temperature that results in the best specificity should be used in subsequent experiments. Fast cycling conditions should not be used for multiplex PCR. Initially, we recommend a 90 second extension time. This time may be further extended to increase yield.

**Colony PCR:** From bacterial colonies use a sterile tip to pick a colony and resuspend into a 50µL reaction as described below. From liquid culture add 5µL of overnight culture to the final mix. Increase initial denaturation time to 10 minutes.

**Direct blood/urine PCR:** Add 2µL mammalian blood or urine to a 50µL reaction as described below.

## Reaction setup

1. Prepare a master mix based on the following table:

Reagent	50µL reaction	Final concentration	Notes
2x PCRBIO HS Taq Mix	25.0µL	1x	
Forward primer (10µM)	2.0µL	400nM	See above for optimal primer design
Reverse primer (10µM)	2.0µL	400nM	
Template DNA	<100ng cDNA, <500ng genomic	variable	See above for template considerations
PCR grade dH <sub>2</sub> O	Up to 50µL final volume		

2. Cycle using conditions based on the following table:

Cycles	Temperature	Time	Notes
1	95°C	1min to 2min	Initial denaturation and enzyme activation. For colony PCR increase denaturation time to 10 minutes
40	95°C	15 seconds	Denaturation
	55°C to 65°C	15 seconds	Anneal
	72°C	1 to 120 seconds	Extension (20 seconds per kb). For multiplex PCR use 90 seconds